

Darouiche RO. Chlorhexidine–alcohol versus povidone–iodine for surgical-site antisepsis. *N Engl J Med* 2010;362:18-26

Webster J, Osborne S. Preoperative bathing or showering with skin antiseptics to prevent surgical site infection (Review). The Cochrane Collaboration. 2009.  
<http://www.thecochranelibrary.com>

Chlebicki M P ET cols. Preoperative chlorhexidine shower or bath for prevention of surgical site infection: A meta-analysis. *AJIC*. 2012.

Eiselt D. Presurgical skin preparation with a novel 2% Chlorhexidine Gluconate Cloth reduces rates of surgical site infection in Orthopaedic surgical patients. *Orthopaedic Nursing*. May/June 2009, 28 (3).

Mangram AJ. Guideline for prevention of surgical site infection, 1999. *Infect Control Hosp Epidemiol* 1999; 20: 250-278

Association Operating Room Nurses. Recommended practices for preoperative patient skin antisepsis. In: *Perioperative Standards and Recommended Practices*. 2009 Edition.

Cambridge, MA. Institute for Healthcare Improvement; 2012 November. (available at [www.ihl.org](http://www.ihl.org))

Langgartner J. Combined skin disinfection with chlorhexidine/propanol and aqueous povidine – iodine reduces bacterial colonisation of central venous catheters. *Intensive Care Med*. 2004;30: 1081-8.

Guzel A. Evaluation of the skin flora after chlorhexidine and povidine-iodine preparation in neurosurgical practice. *Surg Neurol* 2009;71:207-10

Anderson MJ. Efficacy of concurrent application of chlorhexidine gluconate and povidone iodine against six nosocomial pathogens. *Am J Infect Control*. 2010; 38: 826-31.